



Butler University

Digital Commons @ Butler University

Scholarship and Professional Work – COPHS

College of Pharmacy & Health Sciences

1-1998

Comparison of arterial plasma amino acid concentrations in rats fed ad lib oral chow or parenteral nutrition via intravenous or gastric infusion

Jane M. Gervasio
Butler University, jgervasi@butler.edu

Emily B. Hak

Michael C. Storm

M. E. Croft

Lawrence J. Hak

Follow this and additional works at: https://digitalcommons.butler.edu/cophs_papers



Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

Recommended Citation

Gervasio, Jane M.; Hak, Emily B.; Storm, Michael C.; Croft, M. E.; and Hak, Lawrence J., "Comparison of arterial plasma amino acid concentrations in rats fed ad lib oral chow or parenteral nutrition via intravenous or gastric infusion" (1998). *Scholarship and Professional Work – COPHS*. 28.
https://digitalcommons.butler.edu/cophs_papers/28

This Article is brought to you for free and open access by the College of Pharmacy & Health Sciences at Digital Commons @ Butler University. It has been accepted for inclusion in Scholarship and Professional Work – COPHS by an authorized administrator of Digital Commons @ Butler University. For more information, please contact digitalscholarship@butler.edu.

Comparison of arterial plasma amino acid concentrations in rats fed ad lib oral chow or parenteral nutrition via intravenous or gastric infusion

JM Gervasio, EB Hak, MC Storm, ME Croft, LJ Hak

In a rat ischemic acute renal failure model, glycine may be protective to the kidney while lysine and alanine may be nephrotoxic. We have hypothesized that the arterial amino acid plasma pattern, total arterial amino acid concentrations, creatinine clearance, and urine calcium excretion in rats given the same formula via IV or gastrostomy tube continuously would differ. Carotid arterial lines and either an IV or gastrostomy tube were placed in Sprague-Dawley rats who were given parenteral nutrition IV (n = 9), parenteral nutrition via gastrostomy tube (n = 7) or ad lib oral chow (n = 9). After 3 days, arterial blood was drawn during the IV or gastrostomy tube infusion. The ad lib oral chow group was in the absorptive phase. Gastrostomy tube and ad lib oral chow rats gained 5.4 ± 4.6 and 6.8 ± 15 g, respectively, but IV rats lost 12 ± 19 g. Glycine concentrations (nmol/mL) in the IV and gastrostomy tube groups were more than twice the ad lib oral chow group. Alanine and lysine concentrations were not different. Total arterial amino acids were greater in the IV than in the ad lib oral chow group and tended to be greater in the IV group compared to the gastrostomy tube ($p = 0.08$).

	Glycine	Alanine	Lysine	Total amino acids
IV	$721 \pm 209_a$	571 ± 74	612 ± 126	$5239 \pm 439_a$
Gastrostomy tube	$607 \pm 67_a$	556 ± 102	682 ± 110	$4888 \pm 275_a$
Ad lib oral chow	284 ± 57	457 ± 136	536 ± 140	3840 ± 705

P -value ≤ 0.05 = statistical significance; a = different from ad lib oral chow, b = different from gastrostomy tube.

	Urine volume ml	urine calcium mg/dL	Calcium excretion mg/d	Urine creatinine cont'd. mg/dL
IV	$22 \pm 9_{a,b}$	1.3 ± 0.8	11.6 ± 1.2	$17 \pm 12_{a,b}$
Gastrostomy tube	$24 \pm 4_a$	1.6 ± 1.2	11.6 ± 1.6	$17 \pm 7_a$
Ad lib oral chow	$13 \pm 7_b$	7.2 ± 6.8	10.8 ± 1.1	$56 \pm 42_b$

	s creatinine mg/dL	Creatinine clearance ml/min/100g
IV	0.4 ± 0.1	0.28 ± 0.17
Gastrostomy tube	0.5 ± 0.1	0.28 ± 0.14
Ad lib oral chow	0.4 ± 0.1	0.58 ± 0.77

P-value ≤ 0.05 = statistical significance; a = different from ad lib oral chow, b = different from gastrostomy tube.

We conclude that the total amino acid load presented to the kidney is great with IV than with ad lib chow and tends to be greater with IV than with gastrostomy tube infusion of the same parenteral nutrition formula. With parenteral nutrition, arterial glycine concentrations are greater than ad lib oral chow while arterial alanine and lysine concentrations are not different from ad lib oral chow.