



2017

Play It Again! Individualized Music Improves Social Interaction of Women, But Not Men, With Dementia

Emily Farrer
Butler University

Follow this and additional works at: <https://digitalcommons.butler.edu/ugtheses>

 Part of the [Health Psychology Commons](#), [Pain Management Commons](#), and the [Recreational Therapy Commons](#)

Recommended Citation

Farrer, Emily, "Play It Again! Individualized Music Improves Social Interaction of Women, But Not Men, With Dementia" (2017). *Undergraduate Honors Thesis Collection*. 385.
<https://digitalcommons.butler.edu/ugtheses/385>

This Thesis is brought to you for free and open access by the Undergraduate Scholarship at Digital Commons @ Butler University. It has been accepted for inclusion in Undergraduate Honors Thesis Collection by an authorized administrator of Digital Commons @ Butler University. For more information, please contact digitalscholarship@butler.edu.

BUTLER UNIVERSITY HONORS PROGRAM

Honors Thesis Certification

Please type all information in this section:

Applicant Emily Farrer
(Name as it is to appear on diploma)

Thesis title Play It Again! Individualized Music Improves the Social
Interaction of Women, But Not Men, With Dementia

Intended date of commencement May 6, 2017

Read, approved, and signed by:

Thesis adviser(s) Sara Lineweaver 11/28/16
Date

Reader(s) [Signature] 3/15/17
Date
[Signature] 3/17/17
Date

Certified by Rusty Jones 4/13/17
Date
Director, Honors Program

Play It Again! Individualized Music Improves Social Interaction of Women, But Not
Men, With Dementia

A Thesis

Presented to the Department of Psychology

College of Liberal Arts and Sciences

and

The Department of Music

Jordan College of the Arts

and

The Honors Program

Butler University

In Partial Fulfillment

of the Requirements for Graduation Honors

Emily Ann Farrer

3/22/2017

Abstract

Dementia is tragic both for patients with the disease and for their family members. As this neurodegenerative disorder progresses, cognitive functions deteriorate, agitation and aggression increase, and mood worsens (McDermott, Orrell, et al., 2014; Spiro, 2010; Vasionyté & Madison, 2013). Non-pharmacological approaches, such as active music therapy or passive music listening, can be effective in reducing these symptoms. However, no past studies have examined how music listening affects the social interactions of patients with dementia. As such, this study examined whether nursing home residents with dementia are more engaged and interactive if they listen to individually selected music for 15 minutes just prior to a visit with loved ones. Twelve nursing home residents (4 men, 8 women) with dementia ($MMSE M=5.00, SD=7.25$) and their loved ones experienced 5 visits preceded by music and 5 visits not preceded by music, with the exception of one resident who passed away after 3 music and 4 non-music visits. Following each visit, the loved one, blinded to the music vs non-music condition, completed a Social Interaction Questionnaire assessing the resident's alertness, engagement, talkativeness, appropriateness, physical responsiveness, relaxation, happiness, agitation and repetitiveness. Results demonstrated that female residents' social interactions were rated more positively, but male residents' social interactions were rated more negatively, after music than non-music visits. These results support the efficacy of music as an intervention for improving the social interactions of women, but not men, with dementia.

Individualized Music Improves Social Interaction of Women, But Not Men, With
Dementia

Dementia is a neurodegenerative disease characterized by declines in memory, decision-making and judgment, and language skills (American Psychiatric Association, 2000). The corresponding emotional symptoms, including agitation, aggression, depression, and anxiety (Vasionyté & Madison, 2013; Solé, Mercadal-Brotons, Galati & De Castro, 2014), make treatment difficult for both professional caregivers and the patient's family members. According to the World Health Organization (2016), 29 million adults currently suffer from dementia, and the number of diagnosed cases is expected to increase. Thus, dementia is a growing public health concern in today's society.

There is no known cure for dementia, but beyond medications that may slow the progression of the disease, music therapy is one of several types of treatment that has proven to be valuable in combating symptoms (McDermott, Orgeta, Ridder & Orrell, 2014; Spiro, 2010; Vasionyté & Madison, 2013). Music therapy is a relatively new field of study. It was established as a profession in 1950 and is often used as an alternative to pharmacological treatments for dementia patients, since it tends to result in many fewer negative side effects than medications (Vasionyté & Madison, 2013). Solé and colleagues (2014) argue that although pharmacological treatment is important to reduce the effects of dementia, non-pharmacological therapeutic programs are critical to increase mood and improve quality of life in patients suffering from dementing disorders. Music therapy has also proven to increase alertness and enhance the well-being of dementia patients in several studies (McDermott, Orgeta, et al., 2014; Solé et al., 2014).

Music therapy can take two forms: passive (receptive) versus active. In passive therapy participants listen to music without actively contributing, whereas in active therapy the patient is required to participate by singing, playing musical instruments, or clapping (Blackburn & Bradshaw, 2014). Regardless of the type of music therapy, however, individualized music, music that is tailored to the tastes of the individual with dementia, is frequently utilized instead of classical music or calming sounds in order to stimulate reminiscence and emotional responses (Garland, Beer, Eppingstall & O'Connor, 2007; Gerdner & Swanson, 1993; Park, 2010). A study by Gerdner (2000) helped support the importance of utilizing individualized music in music interventions. Family members of patients selected music they believed the patient would enjoy, and music therapists selected relaxing classical music from an anthology. Gerdner divided her sample of 39 patients with Alzheimer's disease into two groups. The first group listened to individualized music for six weeks, followed by two weeks of no music and six weeks of classical music. The second group received the same protocol in reverse, receiving six weeks of classical music first, a two-week washout period, and then six weeks of individualized music. Agitation significantly decreased in both groups within the first 10 minutes of individualized music listening and lasted for approximately 30 minutes following the individualized music. In comparison, classical music did not result in a significant reduction in agitation in either group until 20 minutes into the intervention and improvements lasted only 10 minutes after the classical selections.

Like Gerdner's study (2000), many of the studies that have focused on the benefits of music have examined its impact on agitation. In 1993, Gerdner and Swanson investigated how individualized music listening would affect the agitation of five elderly

patients by having family members select music for their loved ones and having the patients listen to an audio cassette recording of the music for 30 minutes for two weeks. The results indicated a decrease in agitated behaviors in four of the five residents, although the magnitude of the reduction and the length of musical effects varied. Specifically, 47% of agitated behaviors decreased during the intervention, and the improvement was enhanced after the intervention with an 80% decrease in agitated behaviors one-hour post intervention.

A study conducted by Sakamoto, Ando, and Tsutou (2013) supports these findings but also indicates that the improved mood and behavior following passive music listening is a relatively short-term result. Patients with dementia listened to individualized music for half an hour once a week for 10 weeks. The music was selected after in depth interviews with patients and their families. Five minutes before and after each intervention, behavioral and psychological symptoms of dementia were measured. The results indicated that patients were much less agitated after the intervention than before, but when retested two weeks later, the reduction in symptoms was no longer evident.

Based on this literature, an inter-professional group of professors at Butler University, together with several outside consultants, implemented a study called Music First in collaboration with Harrison Terrace Nursing Home (Shiltz et al., 2016). This study used patient-selected music from their late teens to early 20's as a first treatment for agitation in dementia patients. Agitation was measured through nurse and staff reports using the Cohen-Mansfield Agitation Inventory (Cohen-Mansfield, Marx & Rosenthal, 1989). Additional potential effects of music intervention were measured, including its impact on memory, attention, and language skills assessed with the Mini Mental State

Examination (Folstein, Folstein & McHugh, 1975), as well as its effect on hostility, depression, and anxiety, measured using a revised version of the Profile of Mood States (McNair, Lorr & Droppleman, 1992). One hundred and four out of 108 Harrison Terrace residents, all with mild to severe dementia, were enrolled in the study. The results indicated there was a significant interaction between antipsychotic exposure, dementia severity, music listening, and time. Patients who had been receiving a scheduled atypical antipsychotic drug at baseline showed similar levels of agitation to those not on antipsychotics once music listening was added to their care, supporting the hypothesis that individualized music listening may reduce agitation and suggesting it may be an effective adjunct to treatment with medication.

The Music First study was similar to many past studies in its focus on the emotional and behavioral effects of listening to music on dementia patients. However, very little research has assessed the value of the music listening on dementia patients from other people's perspectives. In one such study, McDermott, Orrell, and Ridder (2014) interviewed four separate groups: care home residents with dementia and their families, day hospital clients with dementia, care home staff, and music therapists. McDermott asked the patients questions such as "what does music mean to you?" and she asked families, staff, and therapists questions like "how do you know when music is meaningful to the person?" McDermott found that music helped preserve the self-identity of the patient, decrease the patient's feelings of loneliness, and connect the patient to other people. She also found that both staff and families welcomed the improved mood and increased alertness exhibited by patients after they listened to music.

Taken together, the results of past studies indicate that passive music listening can positively affect dementia patients by increasing their mood and decreasing agitation. Utilizing individualized music rather than general music selections for this intervention is most beneficial, and care providers and family members notice the positive responses residents with dementia have to music. However, the duration of the effects after music listening is typically short-lived. Thus, I have designed the current study to determine whether having residents with dementia listen to music immediately prior to a visit with family members could positively affect the quality of their social interactions during the visit. This would result in a more fulfilling visit for both patients and their loved ones and could lead to the development of more formalized utilization of this inexpensive and simple intervention on a wide scale basis.

Method

Participants

Twelve individuals (4 men, 8 women; age $M=78.67$, $SD=12.09$) participated in this study. All participants were residents at Harrison Terrace Nursing Home, a senior living community in Indianapolis that specializes in memory care, who were diagnosed with mild to severe dementia based on their scores on the Mini-Mental State Examination (MMSE: Folstein, et al., 1975; $M=5.00$, $SD=7.25$). The residents' healthcare proxies and all visiting family members and loved ones gave informed consent before the study began. There were 14 loved ones in total who participated by completing a questionnaire at the conclusion of a visit. Participation was voluntary, and each family received \$50 as a thank you at the conclusion of the study.

Materials

Mini-Mental State Examination (Folstein, et al., 1975). This dementia screening measure provides a baseline to determine the level of dementia severity in each participant. It consists of 30 questions that assess the cognitive abilities of patients including learning and memory, orientation to time and place, language skills, constructional skills, and attention.

Visitation and Listening Log. This log, created for the purposes of this study, recorded information about each visit, including the date, visitor name, length of visit, whether music preceded the visit or not, length of music, and any behaviors exhibited during the music listening. The 11 behaviors documented included, but were not limited to: opening and closing eyes, lifting the head, producing sounds, and moving the body. Please see Appendix A for a copy of this visitation log.

Social Interaction Questionnaire. At the conclusion of each visit, one member of the visiting family completed a 12-item questionnaire, also created for the purposes of this study, about the patient's behavior and emotional state during the visit. The questionnaire included nine behavioral descriptors rated by family members such as how alert, engaged, talkative, and relaxed the resident was during the meeting with their family. Because the exact duration of the positive effects of music therapy is unknown (McDermott, Orgeta, et al., 2014; McDermott, Orrell, et al., 2014; Spaul, Leach & Frampton, 1998), this questionnaire requested ratings for the part of the visit when the patient was most interactive. Please see Appendix B for a copy of this questionnaire.

Personal Playlist. Many participants already had individualized playlists from the Music First study (Shiltz et al., 2016). I created playlists for the remaining participants by

playing hit songs from their late teens to early 20's on an iPad and downloading the songs that generated a behavioral reaction, such as head nodding or snapping. Two families also brought in personal CD's to add to their residents' playlists. At the conclusion of the process, each participant received their own iPod loaded with their personal playlist and a set of headphones.

Procedure

Data collection occurred from May 2015 to October 2015 at Harrison Terrace Nursing Home. I recruited participants through collaboration with Omar Johnson, the Executive Director, Jeannie Keenan RN, the former Director of Nursing Services, and Heather Johnson, LPN, the former Memory Care Specialist. Healthcare proxies of the participants granted consent on the participants' behalf. I administered the Mini-Mental State Examination (Folstein, et al., 1975) to all residents in the study to establish a baseline dementia severity. I quasi-randomly determined whether or not residents listened to music before each visit with family members by flipping a coin while assuring that no more than three consecutive visits were of the same type (music versus non-music visit). Families enrolled in the study notified me an hour in advance of their visit by phone call or email so that I could administer the music or non-music session immediately before the visit without the families' knowledge.

All 12 participants experienced five music and five non-music visits except for one resident who passed away after 3 music visits and 4 non-music visits. For music visits, another researcher or I played selections from the patient's personal playlist for 15 minutes prior to the visit. For non-music visits, we spoke to the resident for approximately five minutes and then returned to the lobby. During all visits, I completed

a Visitation Log in order to monitor the length of the visit, whether music was played or not, and any behavioral reactions it generated. Family members did not know whether or not the residents heard music before the visit. This was to reduce the likelihood that expectancy effects would bias the results of the study. At the conclusion of each visit, one member of the visiting family completed the 12-item Social Interaction Questionnaire before being thanked for their participation. I averaged each residents' scores on each item of the Social Interaction Questionnaire separately for music and non-music visits.

Results

To determine the effect of music versus non-music visits on social interaction, I ran a series of repeated measures ANOVAs for each item on the Social Interaction Questionnaire. I analyzed ratings for each of the nine behavioral descriptors separately to determine which aspects of social interaction were affected by music. Due to the exploratory nature of this research, I treated each item on the Social Interaction Questionnaire as a stand-alone variable. None of the nine analyses yielded a significant difference between the music and non-music conditions, all $F_s(1,11) < 2.07$, all $p_s > .17$.

Because there were no significant results in the primary analysis, I decided to further explore the data by analyzing the effect of gender. This series of exploratory analyses involved nine 2 (Music: music vs. non-music) \times 2 (Gender: male vs. female) mixed-model ANOVAs. Music was a within-subjects factor, whereas gender was a between-subjects factor. Table 1 summarizes the means and standard deviations for male and female residents under each music-listening condition, and Table 2 summarizes the results of these analyses. There were no significant main effects of gender on any of the

nine behaviors. There were also no significant main effects of music, although the music main effect neared significance for appropriateness, $F(1,10)=4.46, p=.061, \eta_p^2=.308$.

Although the main effects failed to reach significance, there was a significant interaction effect between music and gender for alertness ($F(1,10)=5.41, p=.042, \eta_p^2=.351$) and happiness, $F(1,10)=5.16, p=.046, \eta_p^2=.341$. Near significant interactions with moderate effect sizes additionally emerged for engagement ($F(1,10)=3.30, p=.099, \eta_p^2=.248$) and appropriateness, $F(1,10)=3.41, p=.094, \eta_p^2=.254$. (See Table 2.) Figure 1 illustrates the significant and near significant results. Across all of these behaviors, female residents' social interactions were rated more positively, but male residents' social interactions were rated more negatively, after music than non-music visits.

To determine whether residents' reaction to music while they were listening to it predicted the quality of the social interaction that followed, I calculated correlations between the number of observed behaviors during music listening and total scores on the Social Interaction Questionnaire associated with that visit. Table 3 shows the resulting correlation coefficients. Although these correlations were not significant for the first four visits, at the fifth music visit, a significant correlation emerged between residents' behavioral response to music-listening before the visit (e.g., dancing, singing, straightening body/head) and their social interaction during the visit, $r=.65, p<.05$.

Discussion

The primary goal of this study was to determine whether individualized music listening would increase the social interactions of dementia patients as perceived by their visitors. I hypothesized that residents who listened to individualized music prior to visits would be rated as more socially interactive than residents who did not listen to music

prior to visits. The initial analysis indicated there were no significant differences in the nine behavioral descriptors between the music and non-music conditions. However, further analysis of the interaction between music and gender indicated women with dementia demonstrated significantly more happiness and alertness in their social interactions after listening to music than during non-music visits, which supports my hypothesis, but only for female residents. These improvements in the women in my study are consistent with past research that indicates music listening results in positive changes in mood and behavior (Gerdner, 2000; Gerdner & Swanson, 1993; McDermott, Orgeta, et al., 2014; McDermott, Orrell, et al., 2014; Sakamoto, et al., 2013; Vink, Birks, Bruinsma & Scholten, 2013). My results support the use of passive music therapy as a means to enhance social relationships in female dementia residents.

In contrast to the improvements we observed in women, loved ones rated male residents' social interactions more negatively after music than non-music visits, which was unexpected. One possible explanation for this finding is that men showed better social interaction than women in the non-music condition, possibly giving them less room for improvement when music preceded the visit. This may be due to the control condition involving social interactions with the researchers. Even though both researchers made a conscious attempt to keep the interaction short and basic in the control condition, greeting the residents may have produced a positive response and alerted the residents that a family visit was imminent. If this was truer for men than women in the study, this could explain the difference in the findings across the two groups. Additionally, my sample only included four men (half the number of female residents), which likely

limited my ability to document significant differences in the social interactions of men, but not women.

Like past research, my study utilized individualized music in treating dementia patients. Past studies by Garland and colleagues (2007), Gerdner and Swanson (1993), Gerdner (2000), and Ragneskog, Asplund, Kihlgren and Norberg (2001) demonstrate that utilizing individualized music affects dementia patients' mood and behavior more positively than classical or relaxing music. My study applied individualized music listening toward improving social interactions and suggests that this approach can also have a positive effect on nursing home residents' interactions with others. Although past studies have shown that the effects of music listening are temporary (Sakamoto et al., 2013), my study shows that the positive effects of the music last long enough to influence the social interactions that follow.

Because patients with dementia can have varying reactions to the music, I also wanted to determine if their response to music before each visit predicted the quality of their subsequent social interactions. Although there were no significant correlations for the first four visits, at the fifth visit a significant correlation emerged between residents' behavioral responses to the music and the following social interactions. This result suggests that behavioral changes due to music listening may have a cumulative effect over time. A study conducted by Solé and colleagues (2014) supports these findings. They measured quality of life in patients and found that over the course of 12 weekly music therapy sessions there was a significant improvement in median subscale scores for emotional well being from pre-test to post-test. This suggests multiple sessions of music therapy may be necessary to elicit behavioral or emotional changes in patients. Although

other studies also report that the effects of music listening emerge after four to twelve sessions (Garland, Beer, Eppingstall & O'Connor, 2007; Gerdner, 2000; Park, 2010; Sakamoto et al., 2013; Sung, Chang & Lee, 2010), no past studies have examined changes in response to music listening on a trial-by-trial basis. The results of my study indicate there are potential cumulative effects over the course of music listening sessions, which could be examined more in depth in future studies.

While interpreting this data, some limitations should be acknowledged. My sample size was modest, only 12 out of 108 residents at Harrison Terrace participated. I might have been able to detect more changes in social interactions with a larger sample. Additionally, one of the residents passed away during the study, so I did not have data from as many visits for that participant. Due to the small sample size, I also took an exploratory approach to my analyses. Therefore, I did not correct for multiple comparisons when examining each item on the Social Interaction Questionnaire. As such, my Type 1 error rate may be inflated. Future research with larger samples and a more rigorous statistical approach could verify the findings from my study.

Since gender played a primary role in music's effect on happiness and alertness, it is important to acknowledge that there were twice as many women as men enrolled in the study (8 women, 4 men). As previously mentioned, this ratio may have affected the findings that music was an effective treatment for women but not men with dementia. With more men in the sample, I could determine whether the decreased ratings following music listening in men was a robust or chance finding.

A variety of factors may have influenced the accuracy of reported behaviors. The Social Interaction Questionnaire was developed for the purposes of this study and was

therefore not a standardized measurement of social interaction. Also, visitation times varied across visitors so that they completed the questionnaire at different times of the day. Since the behavior of patients with dementia is variable and often dependent on external factors such as meals or sleep, this variation in timing may have affected behavioral ratings (Smith, Gerdner, Hall & Buckwalter, 2004). Individual differences and expectancy effects in visitors who completed the questionnaire may have also affected their behavioral ratings, although I attempted to minimize these by blinding family members to the residents' condition.

The scope of this study was limited since only residents who had frequent and consistent visitors could participate. Future studies could analyze the effects of individualized music listening on all residents displaying agitation, not just those with frequent visitors. Measuring social interaction at various times of the day would help determine when music listening is most beneficial, and monitoring social interactions with other groups of individuals, such as other residents or staff, could evaluate the generalizability of these results. Further research should also assess the optimal length of music listening. For this study residents listened for 15 minutes, but longer listening sessions might have led to larger effects on social interactions. Finally, because some residents displayed irritation after just a few minutes of listening, particularly due to the discomfort of using headphones, sometimes residents listened to music without headphones. Future studies could target the best method to utilize for individualized music listening.

In conclusion, this study is one of the first to examine the effects of gender on individualized music listening so further investigation into this phenomenon is necessary.

However, based on the results of this study, I found support for using individualized music listening to increase the happiness and alertness of women, but not men, with dementia, and these results suggest that music listening may be a simple and inexpensive way to improve the social relationships women with dementia share with others.

References

- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders DSM-IV-TR (Text Revision)*. Washington, DC: American Psychiatric Association.
- Blackburn, R. & Bradshaw, T. (2014). Music therapy for service users with dementia: A critical review of the literature. *Journal of Psychiatric and Mental Health Nursing, 21*(10), 879-888. doi:10.1111/jpm.12165
- Cohen-Mansfield, J., Marx, M. S. & Rosenthal, A. S. (1989). A description of agitation in a nursing home. *Journal of Gerontology: Medical Sciences, 44*(3), 77-84.
- Folstein, M. F., Folstein, S. E. & McHugh, P. R. (1975). "Mini-mental state" a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research, 12*(3), 189-198.
- Garland, K., Beer, E., Eppingstall, B. & O'Connor, D. W. (2007). A comparison of two treatments of agitated behavior in nursing home residents with dementia: Simulated family presence and preferred music. *The American Journal of Geriatric Psychiatry, 15*(6), 514-521. doi:10.1097/01.JGP.0000249388.37080.b4
- Gerdner, L. A. (2000). Effects of individualized versus classical 'relaxation' music on the frequency of agitation in elderly persons with Alzheimer's disease and related disorders. *International Psychogeriatrics, 12*(1), 49-65.
doi:10.1017/S1041610200006190
- Gerdner, L. A. & Swanson, E. A. (1993). Effects of individualized music on confused and agitated elderly patients. *Archives of Psychiatric Nursing, 7*(5), 284-291.
doi:10.1016/0883-9417(93)90006-I

- McDermott, O., Orgeta, V., Ridder, H. M. & Orrell, M. (2014). A preliminary psychometric evaluation of music in dementia assessment scales (MiDAS). *International Psychogeriatrics*, 26(6), 1011-1019.
doi:10.1017/S1041610214000180
- McDermott, O., Orrell, M. & Ridder, H. M. (2014). The importance of music for people with dementia: The perspectives of people with dementia, family carers, staff and music therapists. *Aging and Mental Health*, 18(6), 706-716.
doi:10.1080/13607863.2013.875124
- McNair, D. M., Lorr, M. & Droppleman, L. F. (1992). *POMS manual: profile of mood states*. San Diego, CA: Education and Industrial Testing Service.
- Park, H. (2010). Effect of music on pain for home-dwelling persons with dementia. *Pain Management Nursing*, 11(3), 141-147. doi:10.1016/j.pmn.2009.05.004
- Ragneskog, H., Asplund, K., Kihlgren, M. & Norberg, A. (2001). Individualized music played for agitated patients with dementia: Analysis of video-recorded sessions. *International Journal of Nursing Practice*, 7(3), 146-155. doi:10.1046/j.1440-172X.2001.00254.x
- Sakamoto, M., Ando, H. & Tsutou, A. (2013). Comparing the effects of different individualized music interventions for elderly individuals with severe dementia. *International Psychogeriatrics*, 25(5), 775-784. doi:10.1017/S1041610212002256

- Shiltz, D.L., Lineweaver, T. T., Brimmer, T., Cairns, A., Stone, D., Juett, J., Beer, L., Hay, D. & Plewes, J. (2016, March). "Music first: " *An interprofessional study of the effect of patient-preferred music as a nonpharmacological alternative or adjunct to medications for behavioral and psychological symptoms of dementia*. Poster presented at the meeting of the American Association for Geriatric Psychiatry, Washington, DC.
- Smith, M., Gerdner, L. A., Hall, G. R. & Buckwalter, K. C. (2004). History, development, and future of the progressively lowered stress threshold: A conceptual model for dementia care. *Journal of the American Geriatrics Society*, 52(10), 1755-1760. doi:10.1111/j.1532-5415.2004.52473.x
- Solé, C., Mercadal-Brotons, M., Galati, A. & De Castro, M. (2014). Effects of group music therapy on quality of life, affect, and participation in people with varying levels of dementia. *Journal of Music Therapy*, 51(1), 103-125. doi:10.1093/jmt/thu003
- Spaull, D., Leach, C. & Frampton, I. (1998). An evaluation of the effects of sensory stimulation with people who have dementia. *Behavioural and Cognitive Psychotherapy*, 26(1), 77-86. doi:10.1017/S1352465898000083
- Spiro, N. (2010). Music and dementia: Observing effects and searching for underlying theories. *Aging & Mental Health*, 14(8), 891-899. doi:10.1080/13607863.2010.519328
- Sung, H., Chang, A. & Lee, W. (2010). A preferred music listening intervention to reduce anxiety in older adults with dementia in nursing homes. *Journal of Clinical Nursing*, 19(7-8), 1056-1064. doi:10.1111/j.1365-2702.2009.03016.x

- Ueda, T., Suzukamo, Y., Sato, M. & Izumi, S. I. (2013). Effects of music therapy on behavioral and psychological symptoms of dementia: A systematic review and meta-analysis. *Ageing research reviews*, 12(2), 628-641.
doi:10.1016/j.arr.2013.02.003
- Vasionytė, I. & Madison, G. (2013). Musical intervention for with dementia: A meta-analysis. *Journal of Clinical Nursing*, 22(9-10), 1203-1216.
doi:10.1111/jocn.12166
- Vink, A. C., Birks, J. S., Bruinsma, M. S. & Scholten, R. J. P. M. (2013). Music therapy for people with dementia. *Cochrane Database of Systematic Reviews*, 4.
- WHO (2016). *The World Health report. Executive summary. Mental and neurological disorders*. Retrieved from
http://who.int/whr/1997/media_centre/executive_summary1/en/index14.html on 9.14.2016.

Appendix A

Visitation and Listening Log

Participant ID Number: _____

| Visit # | Date | Visitor Name | Length of Visit | Music Yes or No? | Length of Music | Please circle and note any changes in behavior while listening to music* | | | | |
|---|------|---|-----------------|---|-----------------|--|--|----|-----|------|
| 1 | | | | | | O | L | V | NV | MA |
| In: | | | | | | C | S | SW | SNW | ML D |
| Out: | | | | | | | | | | |
| 2 | | | | | | O | L | V | NV | MA |
| In: | | | | | | C | S | SW | SNW | ML D |
| Out: | | | | | | | | | | |
| 3 | | | | | | O | L | V | NV | MA |
| In: | | | | | | C | S | SW | SNW | ML D |
| Out: | | | | | | | | | | |
| 4 | | | | | | O | L | V | NV | MA |
| In: | | | | | | C | S | SW | SNW | ML D |
| Out: | | | | | | | | | | |
| 5 | | | | | | O | L | V | NV | MA |
| In: | | | | | | C | S | SW | SNW | ML D |
| Out: | | | | | | | | | | |
| 6 | | | | | | O | L | V | NV | MA |
| In: | | | | | | C | S | SW | SNW | ML D |
| Out: | | | | | | | | | | |
| 7 | | | | | | O | L | V | NV | MA |
| In: | | | | | | C | S | SW | SNW | ML D |
| Out: | | | | | | | | | | |
| 8 | | | | | | O | L | V | NV | MA |
| In: | | | | | | C | S | SW | SNW | ML D |
| Out: | | | | | | | | | | |
| 9 | | | | | | O | L | V | NV | MA |
| In: | | | | | | C | S | SW | SNW | ML D |
| Out: | | | | | | | | | | |
| 10 | | | | | | O | L | V | NV | MA |
| In: | | | | | | C | S | SW | SNW | ML D |
| Out: | | | | | | | | | | |
| *Notes Key for behavior changes while listening to music – if more than these are observed, please indicate | | | | | | | | | | |
| Eyes | | Un-coiling | | Sounds | | | Body Movement | | | |
| O = Opens C = Closes | | L = Lifts Head S = Straightens upper torso | | V = Verbal NV = NonVerbal SW = Singing w/ intelligible Words SNW = Singing/humming w/o Words | | | MA = Moves Arms ML = Moves Legs D = Dancing (in any position) | | | |

Appendix B

Social Interaction Questionnaire

Patient Number: _____

Visit Number: _____

Date: _____

1. How long was your visit? Approximately _____ minutes
2. Based upon the visit you just concluded, how would you say the resident's social interactions with you changed over the course of the meeting?
- | | | | | |
|----------|---|----------------------|---|----------|
| worsened | | remained the same | | improved |
| 1 | 2 | 3 | 4 | 5 |
3. Based on the time you just spent with the resident, please indicate the amount of that time the resident was at his or her "best" during your visit.
- Approximately _____ minutes

Based on the time when the resident was *most interactive* during your visit, please circle your answer for the following questions.

4. How alert was the resident?
- | | | | | |
|---------------------|---|-------------------|---|---------------|
| not alert at all | | somewhat alert | | very alert |
| 1 | 2 | 3 | 4 | 5 |
5. How engaged was the resident?
- | | | | | |
|-----------------------|---|---------------------|---|-----------------|
| not engaged at all | | somewhat engaged | | very engaged |
| 1 | 2 | 3 | 4 | 5 |
6. How talkative was the resident?
- | | | | | |
|-------------------------|---|-----------------------|---|-------------------|
| not talkative at all | | somewhat talkative | | very talkative |
| 1 | 2 | 3 | 4 | 5 |

7. How appropriately did the resident respond to comments during conversation?

| | | | | |
|-----------------------------|---|---------------------------|---|-----------------------|
| not appropriately at all | | somewhat appropriately | | very appropriately |
| 1 | 2 | 3 | 4 | 5 |

8. How physically responsive (eye contact, head and body movement) was the resident?

| | | | | |
|--------------------------|---|------------------------|---|--------------------|
| not responsive at all | | somewhat responsive | | very responsive |
| 1 | 2 | 3 | 4 | 5 |

9. How relaxed was the resident?

| | | | | |
|-----------------------|---|---------------------|---|-----------------|
| not relaxed at all | | somewhat relaxed | | very relaxed |
| 1 | 2 | 3 | 4 | 5 |

10. How happy was the resident?

| | | | | |
|---------------------|---|-------------------|---|---------------|
| not happy at all | | somewhat happy | | very happy |
| 1 | 2 | 3 | 4 | 5 |

11. How agitated was the resident?

| | | | | |
|------------------------|---|----------------------|---|------------------|
| not agitated at all | | somewhat agitated | | very agitated |
| 1 | 2 | 3 | 4 | 5 |

12. To what extent did the resident's repetitive behaviors or thoughts interfere with your interaction?

| | | | | |
|---------------|---|----------|---|--------------|
| not at all | | somewhat | | very much |
| 1 | 2 | 3 | 4 | 5 |

Table 1

Social Interaction Ratings for Men and Women with Dementia for Visits Not Preceded by (No Music) or Preceded by (Music) Music

| Behavioral Descriptors | Males | | Females | |
|--------------------------|------------|------------|-------------|------------|
| | No Music | Music | No Music | Music |
| Alert ^a | 3.95 (.30) | 3.60 (.23) | 3.40 (.51) | 3.84 (.72) |
| Engaged ^b | 3.75 (.30) | 3.35 (.25) | 3.20 (.71) | 3.48 (.54) |
| Talkative | 3.20 (.78) | 2.90 (.90) | 2.78 (1.30) | 3.12 (.83) |
| Appropriate ^b | 3.55 (.60) | 3.05 (.84) | 3.30 (.44) | 3.27 (.59) |
| Physically Responsive | 3.70 (.62) | 3.55 (.19) | 3.34 (.72) | 3.45 (.91) |
| Relaxed | 3.85 (.30) | 3.70 (.35) | 3.58 (.57) | 3.91 (.78) |
| Happy ^a | 3.95 (.30) | 3.65 (.19) | 3.18 (.80) | 3.64 (.71) |
| Agitated | 1.90 (.38) | 1.80 (.16) | 1.95 (.75) | 1.85 (.87) |
| Repetitive | 1.85 (.84) | 2.15 (.50) | 2.06 (.40) | 2.04 (.52) |

^aThe Gender x Music interaction was significant ($p < .05$).

^bThe Gender x Music interaction neared significance ($p < .1$).

Table 2

Music Main Effect, Gender Main Effect, and Music x Gender Interaction for Each Item on the Social Interaction Questionnaire

| Behavioral Descriptors | Music Main Effect | | | Gender Main Effect | | | Music x Gender Interaction | | |
|---------------------------|---------------------|----------|------------|---------------------|----------|------------|-------------------------------|----------|------------|
| | <i>F</i> (1, 10) | <i>p</i> | η_p^2 | <i>F</i> (1, 10) | <i>p</i> | η_p^2 | <i>F</i> (1, 10) | <i>p</i> | η_p^2 |
| Alert | .07 | .793 | .01 | .29 | .600 | .03 | 5.41 | .042 | .35 |
| Engaged | .10 | .763 | .01 | .56 | .470 | .05 | 3.30 | .099 | .25 |
| Talkative | .02 | .903 | .00 | .03 | .870 | .00 | 2.65 | .135 | .21 |
| Appropriate | 4.46 | .061 | .31 | .00 | .962 | .00 | 3.41 | .094 | .25 |
| Physically Responsive | .01 | .938 | .00 | .37 | .557 | .04 | .31 | .589 | .03 |
| Relaxed | .28 | .607 | .03 | .01 | .920 | .00 | 1.96 | .192 | .16 |
| Happy | .24 | .632 | .02 | 1.21 | .298 | .11 | 5.16 | .046 | .34 |
| Agitated | .36 | .564 | .03 | .02 | .901 | .00 | .00 | .100 | .00 |
| Repetitive | .46 | .514 | .04 | .04 | .846 | .00 | .60 | .455 | .06 |

Table 3

Correlations Between Behaviors During Music Listening and Scores on the Social Interaction Questionnaire at Each Visit

| Social Interaction | | | | | |
|------------------------|---|----------------|----------------|----------------|----------------|
| Questionnaire Score at | Number of Observed Responses to Music Prior to Each Visit | | | | |
| Each Visit | Visit 1 | Visit 2 | Visit 3 | Visit 4 | Visit 5 |
| Visit 1 | .12 (p=.70) | | | | |
| Visit 2 | | .17 (p=.57) | | | |
| Visit 3 | | | .00 (p=.99) | | |
| Visit 4 | | | | .31 (p=.36) | |
| Visit 5 | | | | | .65 (p=.03) |




Figure 1. Differences in social interactions of dementia patients as measured during visits not preceded by (No Music) or preceded by (Music) music. Behavioral descriptors were measured using the Social Interaction Questionnaire, which used a Likert-type scale ranging from 1 (not at all) to 5 (very). The Gender x Music interaction was significant for alertness ($p=.042$) and happiness ($p=.046$) and neared significance for engagement ($p=.099$) and appropriateness ($p=.094$).