

2015

FOUR COLOR OBSERVATIONS OF 2501 LOHJA

Bin Li

Haibin Zhao

Hao Lu

Xianming Han

Butler University, xhan@butler.edu

Follow this and additional works at: http://digitalcommons.butler.edu/facsch_papers



Part of the [Astrophysics and Astronomy Commons](#)

Recommended Citation

Li, Bin; Zhao, Haibin; Lu, Hao; and Han, Xianming, "FOUR COLOR OBSERVATIONS OF 2501 LOHJA" *Minor Planet Bulletin* / (2015): 190-190.

Available at http://digitalcommons.butler.edu/facsch_papers/740

This Article is brought to you for free and open access by the College of Liberal Arts & Sciences at Digital Commons @ Butler University. It has been accepted for inclusion in Scholarship and Professional Work - LAS by an authorized administrator of Digital Commons @ Butler University. For more information, please contact omacisaa@butler.edu.

FOUR COLOR OBSERVATIONS OF 2501 LOHJA

Bin Li, Haibin Zhao, Hao Lu
Purple Mountain Observatory
Chinese Academy of Sciences
Nanjing 210008. P.R. China
binli@pmo.ac.cn


Xianming L. Han
Dept. of Physics and Astronomy
Butler University
Indianapolis, IN, USA

(Received: 1 April)

Photometric studies of asteroid 2501 Lohja were made between 2014 June 24 and 25 using the Southeastern Association for Research in Astronomy (SARA) Kitt Peak telescope with Bessell B, V, R and I filters. We obtained a synodic period of 3.81 ± 0.01 h, which is consistent with previous values.

All observational data reported here were obtained with the Southeastern Association for Research in Astronomy (SARA) Kitt Peak telescope. This telescope has an aperture of 0.91 m and is located at the Kitt Peak National Observatory near Tucson, Arizona. It has an effective focal ratio of $f/7.5$. When coupled to an Astronomical Research Cameras, Inc. (ARC) CCD camera, the resulting resolution is 0.86 arcsec/pixel (binned 2×2) and the field-of-view (FOV) = $14.6' \times 14.6'$. Bessell BVRI filters were used in turn when taking images. The camera temperature was cooled to -109°C . Image acquisition was done with *DS9*. All images were reduced with master bias, dark, and flat frames. All calibration frames were created using *IDL*. Period analysis was performed using *MPO Canopus*, which incorporates the Fourier analysis algorithm (FALC) developed by Harris (Harris *et al.*, 1989).

We selected 2501 Lohja to accumulate lightcurve data for shape modeling. We also carried out the photometric studies using B, V, R, I filters to detect possible color variations over the surface of the asteroid. Previously reported synodic periods include Higgins (2006, 3.804 h; 2011, 3.80865 h). Husarik (2014) used lightcurve inversion to find the shape, pole, and sidereal period ($P_{\text{Sidereal}} = 3.808348$ h).



We observed the asteroid on 2014 June 24 and 25 and found color indexes of $B-V = 0.94 \pm 0.02$, $V-R = 0.54 \pm 0.01$, and $R-I = 0.382 \pm 0.004$ mag. No significant differences can be seen among these four color lightcurves. We obtained a synodic period $P = 3.81 \pm 0.01$ h, which agrees with the previously determined values.

Acknowledgements

We would like to thank F. Levinson for a generous gift enabling Butler University's membership in the SARA consortium. We would also like to thank the support by the National Natural Science Foundation of China (Grant Nos. 11178025, 11273067 and 10933004), and the Minor Planet Foundation of Purple Mountain Observatory.

References

- Harris, A.W., Young, J.W., Bowell, E., Martin, L.J., Millis, R.L., Poutanen, M., Scaltriti, F., Zappala, V., Schober, H.J., Debehogne, H., Zeigler, K. (1989). "Photoelectric Observations of Asteroids 3, 24, 60, 261, and 863." *Icarus* **77**, 171-186.
- Higgins, D., Pravec, P., Kusnirak, P., Galad, A., Kornos, L., Pray, D., Koff, R.A. (2006). "Asteroid lightcurve analysis at Hunters Hill Observatory and collaborating stations - autumn 2006." *Minor Planet Bulletin* **33**, 89 – 91.
- Higgins, D., Oey, J., Pravec, P. (2011). "Period Determination of Binary Asteroid Targets Observed at Hunters Hill Observatory: May-September 2009." *Minor Planet Bulletin* **38**, 46-49.
- Husarik, M. (2014). "Shapes and spins of asteroids Lohja, Ermolova, and Silver from the observations at the Skalnaté Pleso Observatory." <http://www.helsinki.fi/acm2014/pdf-material/Day-2/Posters-2/HUSARIK-5BE1.pdf>

NEW PHOTOMETRIC OBSERVATIONS OF 128 NEMESIS, 249 ILSE, AND 279 THULE

Frederick Pilcher
4438 Organ Mesa Loop
Las Cruces, NM 88011-8403 USA
fpilcher35@gmail.com

(Received: 20 March Revised: 15 April)

Synodic rotation periods and amplitudes are reported for 128 Nemesis 77.81 ± 0.01 hours, 0.08 ± 0.01 magnitudes; 249 Ilse 84.94 ± 0.01 hours, 0.34 ± 0.02 magnitudes; and 279 Thule, 15.931 ± 0.001 hours, 0.08 ± 0.02 magnitudes.

All observations reported here were made at the Organ Mesa Observatory with a 35 cm Meade LX200 GPS S-C and SBIG STL-1001E CCD, 60 second exposures, unguided. A clear filter was used for 249 Ilse and 279 Thule and R filter for brighter 128 Nemesis. To reduce the large number of data points they have been binned in sets of 3 with maximum time difference 5 minutes for 279 Thule, and in sets of 5 with maximum time difference 10 minutes for 128 Nemesis and 249 Ilse for each of which a total of more than 5000 data points was obtained.