

3-2005

Blood glucose concentration does not correlate with organ failure or outcome in trauma patients receiving enteral nutrition

Jennifer L. Ash
Butler University

Jane M. Gervasio
Butler University, jgervasi@butler.edu

Gary P. Zaloga
Clarian Health Partners

George H. Rodman Jr.
Clarian Health Partners

Follow this and additional works at: http://digitalcommons.butler.edu/cophs_papers

 Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

Recommended Citation

Ash, Jennifer L.; Gervasio, Jane M.; Zaloga, Gary P.; and Rodman Jr., George H., "Blood glucose concentration does not correlate with organ failure or outcome in trauma patients receiving enteral nutrition" (2005). *Scholarship and Professional Work – COPHS*. 24. http://digitalcommons.butler.edu/cophs_papers/24

This Article is brought to you for free and open access by the College of Pharmacy & Health Sciences at Digital Commons @ Butler University. It has been accepted for inclusion in Scholarship and Professional Work – COPHS by an authorized administrator of Digital Commons @ Butler University. For more information, please contact omacisaa@butler.edu.

Blood glucose concentration does not correlate with organ failure or outcome in trauma patients receiving enteral nutrition

Jennifer L. Ash, Jane M. Gervasio, Gary P. Zaloga, George H. Rodman Jr.

Purpose: The objective of this study was to evaluate the relationship between 1) blood glucose concentrations and outcomes and 2) blood glucose concentrations and nutrient intake in critically ill trauma patients receiving enteral nutrition (EN).

Methods: This study is a retrospective chart review assessing the blood glucose concentrations in 120 adult trauma patients receiving EN during the first 7 days in the ICU. The relationship between blood glucose concentrations and organ function (renal, pulmonary, cardiovascular, central nervous system), ventilator days, amount of calories received, ICU LOS, hospital LOS, and number of infections was evaluated using linear regression.

Results: Patients had a mean age of 43.6 ± 18.1 years; 73.3% patients were male; and 8.3% of the population had a diagnosis of diabetes mellitus prior to admission. Mean blood glucose concentrations versus various outcomes are summarized in the following tables:

Table 1. Blood glucose vs. organ failure

Outcome	Renal failure	Respiratory failure	Cardiovascular failure	CNS failure
R ²	0.002	0.023	0.001	0.002

R² = regression coefficient of determination

Table 2. Blood glucose vs. outcomes

Outcome	Caloric intake	Number of infections	Ventilator days	ICU LOS	Hospital LOS
R ²	0.018	0.022	0.010	0.021	0.006

R² = regression coefficient of determination

Conclusions: Blood glucose concentrations were not associated with organ failure, ventilator days, number of infections, ICU LOS, or hospital LOS in critically ill trauma patients receiving EN. In addition, blood glucose concentrations were not affected by the amount of calories administered to trauma patients while in the ICU.